

**BUCCAL, POLAR AND NON-POLAR SPRAY OR CAPSULE CONTAINING
DRUGS FOR TREATING DISORDERS OF THE CENTRAL NERVOUS SYSTEM**

CROSS REFERENCE TO RELATED APPLICATIONS

5 This application is a continuation-in-part of application no. 09/537,118, filed March 29, 2000 which is a continuation-in-part of the U.S. national phase designation of PCT/US97/17899 filed October 1, 1997, the disclosures of which are incorporated by reference herein in their entirety.

10 **BACKGROUND OF THE INVENTION**

 It is known that certain biologically active compounds are better absorbed through the oral mucosa than through other routes of administration, such as through the stomach or intestine. However, formulations suitable for such administration by these latter routes present their own problems. For example, the biologically active compound must be
15 compatible with the other components of the composition such as propellants, solvents, etc. Many such formulations have been proposed. For example, U.S.P. 4,689,233, Dvorsky et al., describes a soft gelatin capsule for the administration of the anti-coronary drug nifedipine dissolved in a mixture of polyether alcohols. U.S.P. 4,755,389, Jones et al., describes a hard gelatin chewable capsule containing nifedipine. A chewable gelatin
20 capsule containing a solution or dispersion of a drug is described in U.S.P. 4,935,243, Borkan et al. U.S.P. 4,919,919, Aouda et al, and U.S.P. 5,370,862, Klokckers-Bethke, describe a nitroglycerin spray for administration to the oral mucosa comprising nitroglycerin, ethanol, and other components. An orally administered pump spray is described by Cholcha in U.S.P. 5,186,925. Aerosol compositions containing a hydrocarbon
25 propellant and a drug for administration to a mucosal surface are described in U.K. 2,082,457, Su, U.S.P. 3,155,574, Silson et al., U.S.P. 5,011,678, Wang et al., and by Parnell in U.S.P. 5,128,132. It should be noted that these references discuss bioavailability of solutions by inhalation rather than through the membranes to which they are administered.

30 **SUMMARY OF THE INVENTION**

 A buccal aerosol spray or soft bite gelatin capsule using a polar or non-polar solvent has now been developed which provides biologically active compounds for rapid absorption through the oral mucosa, resulting in fast onset of effect.

The buccal aerosol spray compositions of the present invention, for transmucosal administration of a pharmacologically active compound soluble in a pharmacologically acceptable non-polar solvent comprise in weight % of total composition: pharmaceutically acceptable propellant 5-80 %, nonpolar solvent 19-85 %, active
5 compound 0.05-50 %, suitably additionally comprising, by weight of total composition a flavoring agent 0.01-10 %. Preferably the composition comprises: propellant 10-70 %, non-polar solvent 25-89.9 %, active compound 0.01-40 %, flavoring agent 1-8 %; most suitably propellant 20-70 %, non-polar solvent 25-74.75 %, active compound 0.25-35 %, flavoring agent 2-7.5 %.

10 The buccal polar aerosol spray compositions of the present invention, for transmucosal administration of a pharmacologically active compound soluble in a pharmacologically acceptable polar solvent are also administrable in aerosol form driven by a propellant. In this case, the composition comprises in weight % of total composition: aqueous polar solvent 10-97 %, active compound 0.1-25 %, suitably additionally
15 comprising, by weight of total composition a flavoring agent 0.05-10 % and propellant: 2 - 10 %. Preferably the composition comprises: polar solvent 20-97 %, active compound 0.1-15%, flavoring agent 0.1-5 % and propellant 2-5 %; most suitably polar solvent 25-97 %, active compound 0.2-25 %, flavoring agent 0.1-2.5 % and propellant 2-4 %.

The buccal pump spray composition of the present invention, *i.e.*, the
20 propellant free composition, for transmucosal administration of a pharmacologically active compound wherein said active compound is soluble in a pharmacologically acceptable non-polar solvent comprises in weight % of total composition: non-polar solvent 30-99.69 %, active compound 0.005-55 %, and suitably additionally, flavoring agent 0.1-10 %.

The buccal polar pump spray compositions of the present invention, *i.e.*, the
25 propellant free composition, for transmucosal administration of a pharmacologically active compound soluble in a pharmacologically acceptable polar solvent comprises in weight % of total composition: aqueous polar solvent 30-99.69 %, active compound 0.001-60 %, suitably additionally comprising, by weight of total composition a flavoring agent 0.1-10 %. Preferably the composition comprises: polar solvent 37-98.58 %, active compound 0.005-55
30 %, flavoring agent 0.5-8 %; most suitably polar solvent 60.9-97.06 %, active compound 0.01-40 %, flavoring agent 0.75-7.5 %.

The soft bite gelatin capsules of the present invention for transmucosal administration of a pharmacologically active compound, at least partially soluble in a

pharmacologically acceptable non-polar solvent, having charged thereto a fill composition
comprise in weight % of total composition: non-polar solvent 4-99.99 %, emulsifier 0-20
%, active compound 0.01-80 %, provided that said fill composition contains less than 10 %
of water, suitably additionally comprising, by weight of the composition: flavoring agent
5 0.01-10 %. Preferably, the soft bite gelatin capsule comprises: non-polar solvent 21.5-
99.975 %, emulsifier 0-15 %, active compound 0.025-70 %, flavoring agent 1-8 %; most
suitably: nonpolar solvent 28.5-97.9 %, emulsifier 0-10 %, active compound 0.1-65.0 %,
flavoring agent 2-6 %.

The soft bite polar gelatin capsules of the present invention for transmucosal
10 administration of a pharmacologically active compound, at least partially soluble in a
pharmacologically acceptable polar solvent, having charged thereto a composition
comprising in weight % of total composition: polar solvent 25-99.89 %, emulsifier 0-20 %,
active compound 0.01-65 %, provided that said composition contains less than 10 % of
water, suitably additionally comprising, by weight of the composition: flavoring agent 01-10
15 %. Preferably, the soft bite gelatin capsule comprises: polar solvent 37-99.95 %, emulsifier
0-15 %, active compound 0.025-55 %, flavoring agent 1-8 %; most suitably: polar solvent
44-96.925 %, emulsifier 0-10 %, active compound 0.075-50 %, flavoring agent 2-6 %.

It is an object of the invention to coat the mucosal membranes either with
extremely fine droplets of spray containing the active compounds or a solution or paste
20 thereof from bite capsules.

It is also an object of the invention to administer to the oral mucosa of a
mammalian in need of same, preferably man, by spray or bite capsule, a predetermined
amount of a biologically active compound by this method or from a soft gelatin capsule.

A further object is a sealed aerosol spray container containing a composition
25 of the non polar or polar aerosol spray formulation, and a metered valve suitable for
releasing from said container a predetermined amount of said composition.

As the propellant evaporates after activation of the aerosol valve, a mist of
fine droplets is formed which contains solvent and active compound.

The propellant is a non-Freon material, preferably a C_{3,8} hydrocarbon of a
30 linear or branched configuration. The propellant should be substantially non-aqueous. The
propellant produces a pressure in the aerosol container such that under expected normal
usage it will produce sufficient pressure to expel the solvent from the container when the
valve is activated but not excessive pressure such as to damage the container or valve seals.

The non-polar solvent is a non-polar hydrocarbon, preferably a C₇₋₁₈ hydrocarbon of a linear or branched configuration, fatty acid esters, and triglycerides, such as miglyol. The solvent must dissolve the active compound and be miscible with the propellant, *i.e.*, solvent and propellant must form a single phase at a temperature of 0-40°C
5 a pressure range of between 1-3 atm.

The polar and non-polar aerosol spray compositions of the invention are intended to be administered from a sealed, pressurized container. Unlike a pump spray, which allows the entry of air into the container after every activation, the aerosol container of the invention is sealed at the time of manufacture. The contents of the container are
10 released by activation of a metered valve, which does not allow entry of atmospheric gasses with each activation. Such containers are commercially available.

A further object is a pump spray container containing a composition of the pump spray formulation, and a metered valve suitable for releasing from said container a predetermined amount of said composition.

15 A further object is a soft gelatin bite capsule containing a composition of as set forth above. The formulation may be in the form of a viscous solution or paste containing the active compounds. Although solutions are preferred, paste fills may also be used where the active compound is not soluble or only partially soluble in the solvent of choice. Where water is used to form part of the paste composition, it should not exceed 10
20 % thereof. (All percentages herein are by weight unless otherwise indicated.)

The polar or non-polar solvent is chosen such that it is compatible with the gelatin shell and the active compound. The solvent preferably dissolves the active compound. However, other components wherein the active compound is not soluble or only slightly soluble may be used and will form a paste fill.

25 Soft gelatin capsules are well known in the art. See, for example, U.S.P. 4,935,243, Borkan et al., for its teaching of such capsules. The capsules of the present invention are intended to be bitten into to release the low viscosity solution or paste therein, which will then coat the buccal mucosa with the active compounds. Typical capsules, which are swallowed whole or bitten and then swallowed, deliver the active compounds to
30 the stomach, which results in significant lag time before maximum blood levels can be achieved or subject the compound to a large first pass effect. Because of the enhanced absorption of the compounds through the oral mucosa and no chance of a first pass effect, use of the bite capsules of the invention will eliminate much of the lag time, resulting in

hastened onset of biological effect. The shell of a soft gelatin capsule of the invention may comprise, for example: gelatin: 50-75 %, glycerin 20-30 %, colorants 0.5-1.5 %, water 5-10 %, and sorbitol 2-10 %.

5 The active compound may include, biologically active peptides, central nervous system active amines, sulfonyl ureas, antibiotics, antifungals, antivirals, sleep inducers, antiasthmatics, bronchial dilators, antiemetics, histamine H-2 receptor antagonists, barbiturates, prostaglandins and neutraceuticals.

The active compounds may also include antihistamines, alkaloids, hormones, benzodiazepines and narcotic analgesics. While not limited thereto, these active compounds
10 are particularly suitable for non-polar pump spray formulation and application.

The active compounds may also include p-FOX (fatty acid oxidation) inhibitors, acetylcholinesterase inhibitors, nerve impulse inhibitors, anti-cholinergics, anti-convulsants, anti-psychotics, anxiolytic agents, dopamine metabolism inhibitors, agents to treat post stroke sequelae, neuroprotectants, agents to treat Alzheimer's disease,
15 neurotransmitters, neurotransmitter agonists, sedatives, agents for treating attention deficit disorder, agents for treating narcolepsy, central adrenergic antagonists, anti-depression agents, agents for treating Parkinson's disease, benzodiazepine antagonists, stimulants, neurotransmitter antagonists, tranquilizers, or a mixture thereof.

20 BRIEF DESCRIPTION OF THE DRAWING

FIG 1. is a schematic diagram showing routes of absorption and processing of pharmacologically active substances in a mammalian system.

DESCRIPTION OF THE PREFERRED EMBODIMENTS

25 The preferred active compounds of the present invention are in an ionized, salt form or as the free base of the pharmaceutically acceptable salts thereof (provided, for the aerosol or pump spray compositions, they are soluble in the spray solvent). These compounds are soluble in the non-polar solvents of the invention at useful concentrations or can be prepared as pastes at useful concentrations. These concentrations may be less than
30 the standard accepted dose for these compounds since there is enhanced absorption of the compounds through the oral mucosa. This aspect of the invention is especially important when there is a large (40-99.99%) first pass effect.

As propellants for the non polar sprays, propane, N-butane, iso-butane, N-pentane, iso-pentane, and neo-pentane, and mixtures thereof may be used. N-butane and iso-butane, as single gases, are the preferred propellants. It is permissible for the propellant to have a water content of no more than 0.2%, typically 0.1-0.2%. All percentages herein are by weight unless otherwise indicated. It is also preferable that the propellant be synthetically produced to minimize the presence of contaminants which are harmful to the active compounds. These contaminants include oxidizing agents, reducing agents, Lewis acids or bases, and water. The concentration of each of these should be less than 0.1 %, except that water may be as high as 0.2%.

Suitable non-polar solvents for the capsules and the non-polar sprays include (C_2 - C_{24}) fatty acid (C_2 - C_6) esters, C_7 - C_{18} hydrocarbon, C_2 - C_6 alkanoyl esters, and the triglycerides of the corresponding acids. When the capsule fill is a paste, other liquid components may be used instead of the above low molecular weight solvents. These include soya oil, corn oil, other vegetable oils.

As solvents for the polar capsules or sprays there may be used low molecular weight polyethyleneglycols (PEG) of 400-1000 Mw (preferably 400-600), low molecular weight (C_2 - C_8) mono and polyols and alcohols of C_7 - C_{18} linear or branch chain hydrocarbons, glycerin may also be present and water may also be used in the sprays, but only in limited amount in the capsules.

It is expected that some glycerin and water used to make the gelatin shell will migrate from the shell to the fill during the curing of the shell. Likewise, there may be some migration of components from the fill to the shell during curing and even throughout the shelf-life of the capsule.

Therefore, the values given herein are for the compositions as prepared, it being within the scope of the invention that minor variations will occur.

The preferred flavoring agents are synthetic or natural oil of peppermint, oil of spearmint, citrus oil, fruit flavors, sweeteners (sugars, aspartame, saccharin, etc.), and combinations thereof.

The active substances include the active compounds selected from the group consisting of cyclosporine, sermorelin, octreotide acetate, calcitonin-salmon, insulin lispro, sumatriptan succinate, clozapine, cyclobenzaprine, dexfenfluramine hydrochloride, glyburide, zidovudine, erythromycin, ciprofloxacin, ondansetron hydrochloride, dimenhydrinate, cimetidine hydrochloride, famotidine, phenytoin sodium, phenytoin,

carboprost thromethamine, carboprost, diphenhydramine hydrochloride, isoproterenol hydrochloride, terbutaline sulfate, terbutaline, theophylline, albuterol sulfate and neutraceuticals, that is to say nutrients with pharmacological action such as but not limited to carnitine, valerian, echinacea, and the like.

5 In another embodiment, the active compound is a p-FOX (fatty acid oxidation) inhibitor, acetylcholinesterase inhibitor, nerve impulse inhibitor, anti-cholinergic, anti-convulsant, anti-psychotic, anxiolytic agent, dopamine metabolism inhibitor, agent to treat post stroke sequelae, neuroprotectant, agent to treat Alzheimer's disease, neurotransmitter, neurotransmitter agonist, sedative, agent for treating attention deficit
10 disorder, agent for treating narcolepsy, central adrenergic antagonist, anti-depression agent, agent for treating Parkinson's disease, benzodiazepine antagonist, stimulant, neurotransmitter antagonist, tranquilizer, or a mixture thereof.

In one embodiment the active compound is a p-FOX inhibitor. A suitable p-FOX inhibitor for use in the buccal sprays of the invention includes, but is not limited to,
15 ranolazine.

In one embodiment the active compound is an acetylcholinesterase inhibitor. Suitable acetylcholinesterase inhibitors for use in the buccal sprays of the invention include, but are not limited to, galantamine, neostigmine, physostigmine, and edrophonium.

In one embodiment the active compound is a nerve impulse inhibitor.
20 Suitable nerve impulse inhibitors for use in the buccal sprays of the invention include, but are not limited to, levobupivacaine, lidocaine, prilocaine, mepivacaine, propofol, rapacuronium bromide, ropivacaine, tubocurarine, atracurium, doxaurium, mivacurium, pancuronium, vecuronium, pipecuronium, and rocuronium.

In one embodiment the active compound is an anti-cholinergic. Suitable
25 anti-cholinergics for use in the buccal sprays of the invention include, but are not limited to, amantadine, ipratropium, oxitropium, and dicycloverine.

In one embodiment the active compound is an anti-convulsant. Suitable anti-convulsants for use in the buccal sprays of the invention include, but are not limited to, acetazolamide, carbamazepine, clonazepam, diazepam, divalproex (valproic acid),
30 ethosuximide, lamotrigine acid, levetiracetam, oxcarbazepine, phenobarbital, phenytoin, pregabalin, primidone, remacemide, trimethadione, topiramate, vigabatrin, and zonisamide.

In one embodiment the active compound is an anti-psychotic. Suitable anti-psychotics for use in the buccal sprays of the invention include, but are not limited to,

amisulpride, aripiprazole bifemelane, bromperidol, clozapine, chlorpromazine, haloperidol, iloperidone loperidone, olanzapine, quetiapine, fluphenazine, fumarate, risperidone, thiothixene, thioridazine, sulpride, and ziprasidone,

5 In one embodiment the active compound is an anxiolytic agent. Suitable anxiolytic agents for use in the buccal sprays of the invention include, but are not limited to, amitryptiline, atracurium, buspirone, chlorzoxazone, clorazepate, cisatracurium, cyclobenzaprine, eperisone, esopiclone, hydroxyzine, mirtazapine, mivacurium, pagoclone, sulperide, zaleplon, and zopiclone.

10 In one embodiment the active compound is a dopamine metabolism inhibitor. Suitable dopamine metabolism inhibitors for use in the buccal sprays of the invention include, but are not limited to, entacapone, lazebemide, selegiline, and tolcapone.

15 In one embodiment the active compound is an agent to treat post stroke sequelae. Suitable agents to treat post stroke sequelae for use in the buccal sprays of the invention include, but are not limited to, glatiramer, interferon beta 1A, interferon beta 1B, estradiol, and progesterone.

In one embodiment the active compound is a neuroprotectant. Suitable neuroprotectants for use in the buccal sprays of the invention include, but are not limited to, donepezil, memantine, nimodipine, riluzole, rivastigmine, tacrine, TAK147, and xaliproden.

20 In one embodiment the active compound is an agent to treat Alzheimer's disease. Suitable agents to treat Alzheimer's disease for use in the buccal sprays of the invention include, but are not limited to, carbidopa, levodopa, tacrine, donezepil, rivastigmine, and galantamine.

25 In one embodiment the active compound is a neurotransmitter. Suitable neurotransmitters for use in the buccal sprays of the invention include, but are not limited to, acetylcholine, serotonin, 5-hydroxytryptamine (5-HT), GABA, glutamate, aspartate, glycine, histamine, epinephrine, norpinephrine, dopamine, adenosine, ATP, and nitric oxide.

30 In one embodiment the active compound is a neurotransmitter agonist. Suitable neurotransmitter agonists for use in the buccal sprays of the invention include, but are not limited to, almotriptan, aniracetam, atomoxetine, benserazide, bromocriptine, bupropion, cabergoline, citalopram, clomipramine, desipramine, diazepam, dihydroergotamine, doxepin duloxetine, eletriptan, escitalopram, fluvoxamine, gabapentin, imipramine, moclobemide, naratriptan, nefazodone, nefiracetam acamprosate, nicergoline,

nortryptiline, paroxetine, pergolide, pramipexole, rizatriptan, ropinirole, sertraline, sibutramine, sumatriptan, tiagabine, trazodone, venlafaxine, and zolmitriptan.

In one embodiment the active compound is a sedative. Suitable sedatives for use in the buccal sprays of the invention include, but are not limited to, dexmedetomidine, eszopiclone, indiplon, zolpidem, and zaleplon.

In one embodiment the active compound is an agent for treating attention deficit disorder. Suitable agents for treating attention deficit disorder for use in the buccal sprays of the invention include, but are not limited to, amphetamine, dextroamphetamine, methylphenidate, and pemoline.

In one embodiment the active compound is an agent for treating narcolepsy. Suitable agents for treating narcolepsy for use in the buccal sprays of the invention include, but are not limited to, modafinil and mazindol.

In one embodiment the active compound is a central adrenergic antagonists. A suitable central adrenergic antagonists for use in the buccal sprays of the invention includes, but is not limited to, mesoridazine.

In one embodiment the active compound is an anti-depression agent. Suitable anti-depression agents for use in the buccal sprays of the invention include, but are not limited to, amitriptyline, amoxapine, bupropion, clomipramine, clomipramine, clorgyline, desipramine, doxepin, fluoxetine, imipramine, isocarboxazid, maprotiline, mirtazapine, nefazodone, nortriptyline, paroxetine, phenelzine, protriptyline, sertraline, tranlycypromine, trazodone, and venlafaxine.

In one embodiment the active compound is an agent for treating Parkinson's disease. Suitable agents for treating Parkinson's disease for use in the buccal sprays of the invention include, but are not limited to, amantadine, bromocriptine, carvidopa, levodopa, pergolide, and selegiline.

In one embodiment the active compound is a benzodiazepine antagonist. A suitable benzodiazepine antagonist for use in the buccal sprays of the invention includes, but is not limited to, flumazenil.

In one embodiment the active compound is a neurotransmitter antagonist. A suitable neurotransmitter antagonist for use in the buccal sprays of the invention includes, but is not limited, to deramciclone.

In one embodiment the active compound is a stimulant. Suitable stimulants for use in the buccal sprays of the invention include, but are not limited to, amphetamine,

dextroamphetamine, dinoprostone, methylphenidate, methylphenidate, modafinil, and pemoline.

In one embodiment the active compound is a tranquilizer. A suitable tranquilizer for use in the buccal sprays of the invention includes, but is not limited to, mesoridazine.

The formulations of the present invention comprise an active compound or a pharmaceutically acceptable salt thereof. The term "pharmaceutically acceptable salts" refers to salts prepared from pharmaceutically acceptable non-toxic acids or bases including organic and inorganic acids or bases.

When an active compound of the present invention is acidic, salts may be prepared from pharmaceutically acceptable non-toxic bases. Salts derived from all stable forms of inorganic bases include aluminum, ammonium, calcium, copper, iron, lithium, magnesium, manganese, potassium, sodium, zinc, etc. Particularly preferred are the ammonium, calcium, magnesium, potassium, and sodium salts. Salts derived from pharmaceutically acceptable organic non-toxic bases include salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion-exchange resins such as arginine, betaine, caffeine, choline, N,N dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, isopropylamine, lysine, methyl-glucosamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purine, theobromine, triethylamine, trimethylamine, tripropylamine, etc.

When an active compound of the present invention is basic, salts may be prepared from pharmaceutically acceptable non-toxic acids. Such acids include acetic, benzenesulfonic, benzoic, camphorsulfonic, citric, ethane-sulfonic, fumaric, gluconic, glutamic, hydrobromic, hydrochloric, isethionic, lactic, maleic, mandelic, methanesulfonic, mucic, nitric, pamoic, pantothenic, phosphoric, succinic, sulfuric, tartaric, p-toluenesulfonic, etc. Particularly preferred are citric, hydrobromic, maleic, phosphoric, sulfuric, and tartaric acids.

In the discussion of methods of treatment herein, reference to the active compounds is meant to also include the pharmaceutically acceptable salts thereof. While certain formulations are set forth herein, the actual amounts to be administered to the mammal or man in need of same are to be determined by the treating physician.

The invention is further defined by reference to the following examples, which are intended to be illustrative and not limiting.

The following are examples of certain classes. All values unless otherwise specified are in weight percent.

5

EXAMPLES

EXAMPLE 1

Biologically active peptides including peptide hormones

A. Cyclosporine lingual spray

5		Amounts	preferred amount	most preferred amount
	cyclosporine	5-50	10-35	15-25
	water	5-20	7.5-50	9.5-12
	ethanol	5-60	7.5-50	10-20
	polyethylene glycol	20-60	30-45	35-40
10	flavors	0.1-5	1-4	2-3

B. Cyclosporine Non-Polar lingual spray

		Amounts	preferred amount	most preferred amount
	cyclosporine	1-50	3-40	5-30
15	Migylol	20	25	30-40
	Polyoxyethylated castor oil	20	25	30-40
	Butane	25-80	30-70	33-50
	flavors	0.1-5	1-4	2-3

20 C. Cyclosporine non-polar bite caosule

		Amounts	preferred amount	most preferred amount
	cyclosporine	1-35	5-25	10-20
	olive oil	25-60	35-55	30-45
25	polyoxyethylated oleic glycerides	25-60	35-55	30-45
	flavors	0.1-5	1-4	2-3

30

D. Cyclosporine bite capsule

	Amounts	preferred amount	most preferred amount
cyclosporine	5-50	10-35	15-25
polyethylene glycol	20-60	30-45	35-40
5 glycerin	5-30	7.5-25	10-20
propylene glycol	5-30	7.5-25	10-20
flavors	0.1-10	1-8	3-6

E. Sermorelin (as the acetate) lingual spray

	Amounts	preferred amount	most preferred
10 sermorelin (as the acetate)	.01-5	.1-3	.2-1.0
mannitol	1-25	5-20	10-15
monobasic sodium phosphate,	0.1-5	1-3	.5-2.5
dibasic sodium phosphate water	0.01-5	.05-3	0.1-0.5
15 ethanol	5-30	7.5-25	9.5-15
polyethylene glycol	20-60	30-45	35-40
propylene glycol	5-25	10-20	12-17
flavors	0.1-5	1-4	2-3

20 F. Octreotide acetate (Sandostatin) lingual spray

	Amounts	preferred amount	most preferred amount
octreotide acetate	0.001-0.5	0.005-0.250	0.01-0.10
acetic acid	1-10	2-8	4-6
sodium acetate	1-10	2-8	4-6
25 sodium chloride	3-30	.5-25	15-20
flavors	0.1-5	0.5-4	2-3
ethanol	5-30	7.5-20	9.5-15
water	15-95	35-90	65-85
flavors	0.1-5	1-4	2-3

30

G. Calcitonin-salmon lingual spray

		Amounts	preferred amount	most preferred amount
	calcitonin-salmon	0.001-5	0.005-2	01-1.5
	ethanol	2-15	3-10	7-9.5
5	water	30-95	50-90	60-80
	polyethylene glycol	2-15	3-10	7-9.5
	sodium chloride	2.5-20	5-15	10-12.5
	flavors	0.1-5	1-4	2-3

10 H. Insulin lispro, lingual spray

		Amounts	preferred amount	most preferred amount
	insulin	20-60	4-55	5-50
	glycerin	0.1-10	0.25-5	0.1-1.5
	dibasic sodium phosphate	1-15	2.5-10	4-8
15	m-cresol,	1-25	5-25	7.5-12.5
	zinc oxide	0.01-0.25	.05-0.15	0.075-0.10
	m-cresol	0.1-1	0.2-0.8	0.4-0.6
	phenol	trace amounts	trace amounts	trace amounts
	ethanol	5-20	7.5-15	9-12
20	water	30-90	40-80	50-75
	propylene glycol	5-20	7.5-15	9-12
	flavors	0.1-5	0.5-3	0.75-2

adjust pH to 7.0-7.8 with HCl or NaOH

25

EXAMPLE 2

CNS active amines and their salts: including but not limited to tricyclic amines, GABA analogues, thiazides, phenothiazine derivatives, serotonin antagonists and serotonin reuptake inhibitors

5 A. Sumatriptan succinate lingual spray

	Amounts	preferred amount	most preferred amount
sumatriptan succinate	0.5-30	1-20	10-15
ethanol	5-60	7.5-50	10-20
propylene glycol	5-30	7.5-20	10-15
10 polyethylene glycol	0-60	30-45	35-40
water	5-30	7.5-20	10-15
flavors	0.1-5	1-4	2-3

B. Sumatriptan succinate bite capsule

	Amounts	preferred amount	most preferred amount
sumatriptan succinate	0.01-5	0.05-3.5	0.075-1.75
polyethylene glycol	25-70	30-60	35-50
glycerin	25-70	30-60	35-50
flavors	0.1-10	1-8	3-6

20

C. Clozepine lingual spray

	Amounts	preferred amount	most preferred amount
clozepine	0.5-30	1-20	10-15
ethanol	5-60	7.5-50	10-20
25 propylene glycol	5-30	7.5-20	10-15
polyethylene glycol	0-60	30-45	35-40
water	5-30	7.5-20	10-15
flavors	0.1-5	1-4	2-3

30

D. Clozepine non-polar lingual spray with propellant

		Amounts	preferred amount	most preferred amount
	clozepine	0.5-30	1-20	10-15
5	Migylol	20-85	25-70	30-40
	Butanol	5-80	30-75	60-70
	flavors	0.1-5	1-4	2-3

E. Clozepine non-polar lingual spray without propellant

		Amounts	preferred amount	most preferred amount
10	clozepine	0.5-30	1-20	10-15
	Migylol	70-99.5	80-99	85-90
	flavors	0.1-5	1-4	2-3

15 F. Cyclobenzaprine non-polar lingual spray

		Amounts	preferred amount	most preferred amount
	cyclobenzaprine (base)	0.5-30	1-20	10-15
	Migylol	20-85	25-70	30-40
	Iso-butane	15-80	30-75	60-70
20	flavors	0.1-5	1-4	2-3

G. Dexfenfluramine hydrochloride lingual spray

		Amounts	preferred amount	most preferred amount
25	dexfenfluramine Hcl	5-30	7.5-20	10-15
	ethanol	5-60	7.5-50	10-20
	propylene glycol	5-30	7.5-20	10-15
	polyethylene glycol	0-60	30-45	35-40
	water	5-30	7.5-20	10-15
30	flavors	0.1-5	1-4	2-3

EXAMPLE 3

Sulfonylureas

A. Glyburide lingual spray

		Amounts	preferred amount	most preferred amount
5	glyburide	0.25-25	0.5-20	0.75-15
	ethanol	5-60	-7.5-50	10-20
	propylene glycol	5-30	7.5-20	10-15
	polyethylene glycol	0-60	30-45	35-40
	water	2.5-30	5-20	6-15
10	flavors	0.1-5	1-4	2-3

B. Glyburide non-polar bite capsule

		Amounts	preferred amount	most preferred amount
	glyburide	0.01-10	0.025-7.5	0.1-4
15	olive oil	30-60	35-55	30-50
	polyoxyethylated oleic glycerides	30-60	35-55	30-50
	flavors	0.1-5	1-4	2-3

20

EXAMPLE 4

Antibiotics anti-fungals and anti-virals

A. Zidovudine [formerly called azidothymidine (AZT) (Retrovir)] non-polar lingual spray

5		Amounts	preferred amount	most preferred amount
	zidovudine	10-50	15-40	25-35
	Soya oil	20-85	25-70	30-40
	Butane	15-80	30-75	60-70
	flavors	0.1-5	1-4	2-3

10

B. Erythromycin bite capsule bite capsule

		Amounts	preferred amount	most preferred amount
	erythromycin	25-65	30-50	35-45
	polyoxyethylene glycol	5-70	30-60	45-55
15	glycerin	5-20	7.5-15	10-12.5
	flavors	1-10	2-8	3-6

C. Ciprofloxacin hydrochloride bite capsule

		Amounts	preferred amount	most preferred amount
20	ciprofloxacin hydrochloride	25-65	35-55	40-50
	glycerin	5-20	7.5-15	10-12.5
	polyethylene glycol	120-75	30-65	40-60
	flavors	1-10	2-8	3-6

25 D. zidovudine [formerly called azidothymidine (AZT) (Retrovir)] lingual spray

		Amounts	preferred amount	most preferred amount
	zidovudine	10-50	15-40	25-35
	water	30-80	40-75	45-70
	ethanol	5-20	7.5-15	9.5-12.5
30	polyethylene glycol	5-20	7.5-15	9.5-12.5
	flavors	0.1-5	1-4	2-3

EXAMPLE 5

Anti-emetics

A. Ondansetron hydrochloride lingual spray

		Amounts	preferred amount	most preferred amount
5	ondansetron hydrochloride	1-25	2-20	2.5-15
	citric acid monohydrate	1-10	2-8	2.5-5
	sodium citrate dihydrate	0.5-5	1-4	1.25-2.5
	water	1-90	5-85	10-75
	ethanol	5-30	7.5-20	9.5-15
10	propylene glycol	5-30	7.5-20	9.5-15
	polyethylene glycol	5-30	7.5-20	9.5-15
	flavors	1-10	3-8	5-7.5

B. Dimenhydrinate bite capsule

		Amounts	preferred amount	most preferred amount
15	dimenhydrinate	0.5-30	2-25	3-15
	glycerin	5-20	7.5-15	10-1 2.5
	polyethylene glycol	45-95	50-90	55-85
	flavors	1-10	2-8	3-6

20

C. Dimenhydrinate polar lingual spray

		Amounts	preferred amount	most preferred amount
	dimenhydrinate	3-50	4-40	5-35
	water	5-90	10-80	15-75
25	ethanol	1-80	3-50	5-10
	polyethylene glycol	1-80	3-50	5-15
	sorbitol	0.1-5	0.2-40	0.4-1.0
	aspartame	0.01-0.5	0.02-0.4	0.04-0.1
	flavors	0.1-5	1-4	2-3

30

EXAMPLE 6

Histamine H-2 receptor antagonists

A. Cimetidine hydrochloride bite capsule

		Amounts	preferred amount	most preferred amount
5	cimetidine HCl	10-60	15-55	25-50
	glycerin	5-20	7.5-15	10-12.5
	polyethylene glycol	20-90	25-85	30-75
	flavors	1-10	2-8	3-6

10 B. Famotidine lingual spray

		Amounts	preferred amount	most preferred amount
	famotidine	1-35	5-30	7-20
	water	2.5-25	3-20	5-10
	L-aspartic acid	0.1-20	1-15	5-10
15	polyethylene glycol	20-97	30-95	50-85
	flavors	0.1-10	1-7.5	2-5

C. Famotidine non-polar lingual spray

		Amounts	preferred amount	most preferred amount
20	famotidine	1-35	5-30	7-20
	Soya oil	10-50	15-40	15-20
	Butane1	5-80	30-75	45-70
	polyoxyethylated oleic glycerides	10-50	15-40	15-20
25	flavors	0.1-5	1-4	2-3

EXAMPLE 7

Barbiturates

A. Phenytoin sodium lingual spray

		Amounts	preferred amount	most preferred amount
5	phenytoin sodium	10-60	15-55	20-40
	water	2.5-25	3-20	5-10
	ethanol	5-30	7.5-20	9.5-15
	propylene glycol	5-30	7.5-20	9.5-15
	polyethylene glycol	5-30	7.5-20	9.5-15
10	flavors	1-10	3-8	5-7.5

B. Phenytoin non-polar lingual spray

		Amounts	preferred amount	most preferred amount
	phenytoin	5-45	10-40	15-35
15	migylol	10-50	15-40	15-20
	Butane	15-80	30-75	60-70
	polyoxyethylated oleic glycerides	10-50	15-40	15-20
	flavors	0.1-10	1-8	5-7.5

20

25

30

EXAMPLE 8

Prostaglandins

A. Carboprost thromethamine lingual spray

		Amounts	preferred amount	most preferred amount
5	carboprost thromethamine	0.05-5	0.1-3	0.25-2.5
	water	50-95	60-80	65-75
	ethanol	5-20	7.5-15	9.5-12.5
	polyethylene glycol	5-20	7.5-15	9.5-12.5
	sodium chloride	1-20	3-15	4-8
10	flavors	0.1-5	1-4	2-3

pH is adjusted with sodium hydroxide and/or hydrochloric acid

B. Carboprost non-polar lingual spray

		Amounts	preferred amount	most preferred amount
15	carboprost	0.05-5	0.1-3	0.25-2.5
	migylol	25-50	30-45	35-40
	Butane	5-60	10-50	20-35
20	polyoxyethylated oleic glycerides	25-50	30-45	35-40
	flavors	0.1-10	1-8	5-7.5

EXAMPLE 9

Neutraceuticals

A. Carnitine as bite capsule (contents are a paste)

		Amounts	preferred amount	most preferred amount
5	carnitine fumarate	6-80	30-70	45-65
	soya oil	7.5-50	10-40	12.5-35
	soya lecithin	0.001-1.0	0.005-0.5	.01-0.1
	Soya fats	7.5-50	10-40	12.5-35
	flavors	1-10	2-8	3-6

10

B. Valerian as lingual spray

		Amounts	preferred amount	most preferred amount
	valerian extract	0.1-10	0.2-7	0.25-5
	water	50-95	60-80	65-75
15	ethanol	5-20	7.5-15	9.5-12.5
	polyethylene glycol	5-20	7.5-15	9.5-12.5
	flavors	1-10	2-8	3-6

C. Echinacea as bite capsule

		Amounts	preferred amount	most preferred amount
20	echinacea extract	30-85	40-75	45-55
	soya oil	7.5-50	10-40	12.5-35
	soya lecithin	0.001-1.0	0.005-0.5	.01-0.1
	Soya fats	7.5-50	10-40	12.5-35
25	flavors	1-10	2-8	3-6

30

D. Mixtures of ingredients

		Amounts	preferred amount	most preferred amount
	magnesium oxide	15-40	20-35	25-30
	chromium picolinate	0.01-1.0	0.02-0.5	.025-0.75
5	folic acid	.025-3.0	0.05-2.0	0.25-0.5
	vitamin B-12	0.01-1.0	0.02-0.5	.025-0.75
	vitamin E	15-40	20-35	25-30
	Soya oil	10-40	12.5-35	15-20
	soya lecithin	0.1-5	0.2-4	0.5-1.5
10	soya fat	10-40	15-35	17.5-20

EXAMPLE 10

Sleep Inducers (also CNS active amine)

A. Diphenhydramine hydrochloride lingual spray

		Amounts	preferred amount	most preferred amount
5	diphenhydramine	3-50.	4-40	5-35
	HCl water	5-90	10-80	50-75
	ethanol	1-80	3-50	5-10
	polyethylene glycol	1-80	3-50	5-15
10	Sorbitol	0.1-5	0.2-4	0.4-1.0
	aspartame	0.01-0.5	0.02-0.4	0.04-0.1
	flavors	0.1-5	1-4	2-3

EXAMPLE 11

Anti-Asthmatics-Bronchodilators

A. Isoproterenol Hydrochloride as polar lingual spray

		Amounts	preferred amount	most preferred amount
5	isoproterenol Hydrochloride	0.1-10	0.2-7.5	0.5-6
	water	5-90	10-80	50-75
	ethanol	1-80	3-50	5-10
	polyethylene glycol	1-80	3-50	5-15
	Sorbitol	0.1-5	0.2-4	0.4-1.0
10	aspartame	0.01-0.5	0.02-0.4	0.04-0.1
	flavors	0.1-5	1-4	2-3

B. Terbutaline sulfate as polar lingual spray

		Amounts	preferred amount	most preferred amount
15	terbutaline sulfate	0.1-10	0.2-7.5	0.5-6
	water	5-90	10-80	50-75
	ethanol	1-10	2-8	2.5-5
	Sorbitol	0.1-5	0.2-4	0.4-1.0
	aspartame	0.01-0.5	0.02-0.4	0.04-0.1
20	flavors	0.1-5	1-4	2-3

C. Terbutaline as non-polar lingual spray

		Amounts	preferred amount	most preferred amount
	terbutaline	0.1-10	0.2-7.5	0.5-6
25	migylol	25-50	30-45	35-40
	isobutane	5-60	10-50	20-35
	polyoxyethylated oleic glycerides	25-50	30-45	35-40
	flavors	0.1-10	1-8	5-7.5
30				

D. Theophylline polar bite capsule

		Amounts	preferred amount	most preferred amount
	theophylline	5-50	10-40	15-30
5	polyethylene glycol	20-60	25-50	30-40
	glycerin	25-50	35-45	30-40
	propylene glycol	25-50	35-45	30-40
	flavors	0.1-5	1-4	2-3

10 E. Albuterol sulfate as polar lingual spray

		Amounts	preferred amount	most preferred amount
	albuterol sulfate	0.1-10	0.2-7.5	0.5-6
	water	5-90	10-80	50-75
	ethanol	1-10	2-8	2.5-5
15	Sorbitol	0.1-5	0.2-4	0.4-1.0
	aspartame	0.01-0.5	0.02-0.4	0.04-0.1
	flavors	0.1-5	1-4	2-3

Example 12

Polar solvent formulations using a propellant:

A. Sulfonylurea

		Amount	Preferred Amount	Most-Preferred Amount
5	glyburide	0.1-25%	0.5-15%	0.6-10%
	Ethanol	40-99%	60-97%	70-97%
	Water	0.01-5%	0.1-4%	0.2-2%
	Flavors	0.05-10%	0.1-5%	0.1-2.5%
	Propellant	2-10%	3-5%	3-4%

10

B. Prostaglandin E (vasodilator)

		Amount	Preferred Amount	Most-Preferred Amount
	prostaglandin E ₁	0.01-10%	0.1-5%	0.2-3%
	Ethanol	10-90%	20-75%	25-50%
15	Propylene glycol	1-90%	5-80%	10-75%
	Water	0.01-5%	0.1-4%	0.2-2%
	Flavors	0.05-10%	0.1-5%	0.1-2.5%
	Propellant	2-10%	3-5%	3-4%

20

C. Promethazine (antiemetic, sleep inducer, and CNS active amine)

		Amount	Preferred Amount	Most-Preferred Amount
	promethazine	1-25%	3-15%	5-12%
	Ethanol	10-90%	20-75%	25-50%
	Propylene glycol	1-90%	5-80%	10-75%
25	Water	0.01-5%	0.1-4%	0.2-2%
	Flavors	0.05-10%	0.1-5%	0.1-2.5%
	Propellant	2-10%	3-5%	3-4%

30

D. Meclizine

	Amount	Preferred Amount	Most-Preferred Amount
meclizine	1-25%	3-15%	5-12%
Ethanol	1-15%	2-10%	3-6
5 Propylene glycol	20-98%	5-90%	10-85%
Water	0.01-5%	0.1-4%	0.2-2%
Flavors	0.05-10%	0.1-5%	0.1-2.5%
Propellant	2-10%	3-5%	3-4%

10